

Epicyclic Frequencies and Resonant Phenomena Near Black Holes: The Current Status

ALIKRAM N. ALIEV*

Feza Gürsey Institute, P.K. 6 Çengelköy, 34684 Istanbul, Turkey

We briefly discuss the theory of epicyclic motion near rotating black holes and its relation to the origin of high frequency Quasi-Periodic Oscillations (QPOs) seen in many cases of accreting black hole binaries. We also point out some new frequency ratios predicted by the theory.

In the framework of general relativity the successive theory of epicyclic motion around rotating black holes was first developed in Refs.^{1,2}(see also a review paper³). As is known, the rotating black holes are uniquely described by the famous Kerr solution

$$ds^2 = -\frac{\Delta}{\Sigma} (dt - a \sin^2 \theta d\phi)^2 + \Sigma \left(\frac{dr^2}{\Delta} + d\theta^2 \right) + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + a^2) d\phi]^2, \quad (1)$$

where the metric functions

$$\Delta = r^2 + a^2 - 2Mr, \quad \Sigma = r^2 + a^2 \cos^2 \theta.$$

The equation of motion for a test particle of mass μ and charge e has the form

$$\frac{d^2 x^\mu}{ds^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} = \frac{e}{\mu} F^\mu{}_\nu \frac{dx^\nu}{ds}. \quad (2)$$

From the symmetry properties of the metric (1) it follows that a cyclic motion of the test particle is possible in the equatorial plane $\theta = \pi/2$. For the cyclic motion we have $z^\mu(s) = u^0 s \{1, 0, 0, \omega_0\}$, where ω_0 is the angular velocity of the motion.

To describe the epicyclic motion of the test particle we employ a perturbative expansion about the cyclic orbits, i.e. we assume that $x^\mu(s) = z^\mu(s) + \xi^\mu(s)$. Substituting this into equation (2) and adding also the external force $f^\mu = f^\mu{}_\nu u^\nu (u^0)^{-1}$, we arrive at the equation

$$\frac{d^2 \xi^\mu}{dt^2} + \gamma_\alpha^\mu \frac{d\xi^\alpha}{dt} + \xi^a \partial_a U^\mu = \frac{e}{\mu u^0} f^\mu + \mathcal{N}^\mu \left(\xi, \frac{d\xi}{dt} \right), \quad a = 1, 2 \equiv r, \theta \quad (3)$$

*aliev@gursey.gov.tr

where $\mathcal{N}^\mu$ stands for the non-linear terms, the quantities γ_α^μ and $\partial_a U^\mu$ must be taken on a cyclic orbit $r = r_0$, $\theta = \pi/2$ and

$$\gamma_\alpha^\mu = 2\Gamma_{\alpha\beta}^\mu u^\beta (u^0)^{-1} - \frac{e}{\mu u^0} F_\alpha^\mu, \quad U^\mu = \frac{1}{2} \left[\gamma_\alpha^\mu u^\alpha (u^0)^{-1} - \frac{e}{\mu u^0} F_\alpha^\mu u^\alpha (u^0)^{-1} \right] \quad (4)$$

Performing an appropriate integration in (3) and restricting ourselves to the first order terms in ξ we obtain the set of equations

$$\frac{d\xi^A}{dt} + \gamma_1^A \xi^r = \frac{e}{\mu u^0} \int f^A dt, \quad A = 0, 3 \equiv t, \phi \quad (5)$$

$$\frac{d^2 \xi^r}{dt^2} + \omega_r^2 \xi^r = \frac{e}{\mu u^0} \left(f^r - \gamma_A^1 \int f^A dt \right), \quad (6)$$

$$\frac{d^2 \xi^\theta}{dt^2} + \omega_\theta^2 \xi^\theta = \frac{e}{\mu u^0} f^\theta. \quad (7)$$

We see that in the absence of the external force the epicyclic motion is governed by the two decoupled oscillatory-type equations. In other words, it amounts to free radial and vertical oscillations with the associated frequencies

$$\omega_r = \left(\frac{\partial U^r}{\partial r} - \gamma_A^1 \gamma_1^A \right)^{1/2}, \quad \omega_\theta = \left(\frac{\partial U^\theta}{\partial \theta} \right)^{1/2}. \quad (8)$$

The explicit expressions for these frequencies were first found in Ref.¹ in the most general case of black holes with an electric charge or with an external uniform magnetic field. Further, the epicyclic frequencies were used to explore the maser effect near black holes² as well as the effects of periastron precession and the Lense-Thirring drag in the Schwarzschild and Kerr fields threaded by a cosmic string.⁴ For a "pure" Kerr field the frequencies are^{1,2}

$$\omega_r^2 = \omega_0^2 \left(1 - \frac{6M}{r} - \frac{3a^2}{r^2} \pm 8a\omega_K \right), \quad \omega_\theta^2 = \omega_0^2 \left(1 + \frac{3a^2}{r^2} \mp 4a\omega_K \right), \quad (9)$$

where $\omega_0 = \pm \omega_K / (1 \pm a\omega_K)$, $\omega_K = \sqrt{M/r^3}$ is the Kepler frequency and the two signs refer to direct and retrograde motions. Later, in 1987 our expression for the frequency of radial oscillations was re-derived in Ref.⁵ while the formula for the frequency of vertical oscillations re-appeared⁶ only in 1990.

In Ref.¹ we have also pointed out that when taking into account the non-linear terms on the right-hand-side of Eq. (3) a coupling between the oscillations occurs. If the frequencies ω_r and ω_θ are in the rational relation $k_r \omega_r = k_\theta \omega_\theta$, where k_r and k_θ are integers, the resonances must take place. We have plotted the positions of low-order ($k = |k_r| + |k_\theta|$) resonances: $k = 3, 4, 5$, or $(2 : 1, 3 : 1, 3 : 2)$. In recent developments, this theory has been extensively used to explain the frequency ratios and the origin of high frequency Quasi-Periodic Oscillations (QPOs) observed in many cases of accreting black hole or neutron star binaries (see papers^{7,8} and references therein). In some cases, even our plots for the positions of the non-linear resonances were exactly reproduced (see for instance, Ref.⁹).

Our model also admits the appearance of the forced resonances under a perturbing force. Solving Eqs.(5)-(7) for Fourier transforms we obtain

$$\xi^A(\omega, m) = -\frac{i}{\omega_m} \gamma_1^A \xi^r(\omega, m) - \frac{e}{\mu u^0} \frac{f^A(\omega, m)}{\omega_m^2}, \quad (10)$$

$$\xi^r(\omega, m) = \frac{e}{\mu u^0} \frac{f^r(\omega, m) - \frac{i}{\omega_m} \gamma_A^1 f^A(\omega, m)}{\omega_r^2 - \omega_m^2}, \quad (11)$$

$$\xi^\theta(\omega, m) = \frac{e}{\mu u^0} \frac{f^\theta(\omega, m)}{\omega_\theta^2 - \omega_m^2}, \quad \omega_m = \omega - m\omega_0. \quad (12)$$

We see that at the frequencies $\omega = m\omega_0$, $\omega = \Omega_r^\pm = m\omega_0 \pm \omega_r$, $\omega = \Omega_\theta^\pm = m\omega_0 \pm \omega_\theta$ the system exhibits the resonant behaviour.² The similar type of the resonant frequencies was discussed by Kato¹⁰ within a model for kHz QPOs resonantly induced in warped disks of X-ray binaries. In the latest developments, we have performed a complete numerical analysis of the frequency ratios pertaining to the orbits where the frequency of the radial oscillations approaches its maximum value. Some results for direct orbits and for masses $\sim 10M_\odot$ are given in the Table.

a/M	r_{max}/M	ν_0 (Hz)	ν_r (Hz)	ν_θ (Hz)	ν_θ/ν_r	ν_0/ν_r	ν_0/ν_θ
0.00	8.00	141.42	70.71	141.42	2.00	2.00	1.00
0.10	7.58	152.60	75.73	151.17	2.00	2.02	1.01
0.20	7.15	165.68	81.52	162.37	1.99	2.03	1.02
0.30	6.70	181.26	88.27	175.45	1.99	2.05	1.03
0.40	6.24	199.97	96.29	190.74	1.98	2.08	1.05
0.50	5.76	223.29	105.99	209.23	1.97	2.11	1.07
0.60	5.26	252.90	118.03	231.78	1.96	2.14	1.09
0.70	4.71	292.71	133.53	260.59	1.95	2.19	1.12
0.80	4.11	349.83	154.57	298.90	1.93	2.26	1.17
0.90	3.42	442.93	185.95	353.95	1.90	2.38	1.25
0.99	2.45	662.11	243.45	447.52	1.84	2.72	1.48

Acknowledgements: The author thanks the Scientific and Technological Research Council of Turkey (TÜBİTAK) for partial support under the Project 105T437.

References

1. A. N. Aliev and D. V. Gal'tsov, Gen. Relat. Gravit. **13**, 899 (1981).
2. A. N. Aliev, D. V. Gal'tsov and V. I. Petukhov, Astr. Space Sci. **124**, 137 (1986).
3. A. N. Aliev and D. V. Gal'tsov, Sov. Phys. Usp. **32**(1), 75 (1989).
4. A. N. Aliev and D. V. Gal'tsov, Sov. Astron. Lett. **14**(1), 48 (1988).
5. A. Okazaki, S. Kato and J. Fukue, Publ. Astron. Soc. Japan **39**, 457 (1987).
6. S. Kato, Publ. Astron. Soc. Japan **42**, 99 (1990).
7. A. Merloni, M. Vietri, L. Stella and D. Bini, Mon. Not. R. Ast. Soc. **304**, 155 (1999).
8. G. Török, M. Abramowicz, W. Kluźniak and Z. Stuchlik, Ast.& Ap. **436**, 1 (2005).
9. M. A. Abramowicz and W. Kluźniak, astro-ph/0312396
10. S. Kato, astro-ph/0506595.